

Transverse doubled knots and clasp disks

Ichiro Torisu

*Naruto University of Education, 748, Nakajima, Takashima, Naruto-cho, Naruto-shi, 772-8502 Japan
e-mail: torisu@naruto-u.ac.jp*

Date: May 2, 2005.

Abstract

We analyze transverse doubled knots in the standard contact 3-space by using spanned clasp disks. As applications, we will estimate their self-linking number and furthermore we will show that in many cases, transverse twist knots with the maximal self-linking number are unique up to transverse isotopy.

1 Introduction

A *contact structure* on a 3-manifold is a completely non-integrable 2-plane field on it. Let ξ_0 be the standard contact structure on 3-space $R^3 = \{(x, y, z)\}$, that is, a 2-plane field on R^3 defined by the kernel of the 1-form $dz - ydx$. A *transverse knot* is an oriented knot which is everywhere transverse to the contact structure. Throughout, we always assume a transverse knot is positively oriented to ξ_0 . A *Legendrian knot* is an oriented knot which is everywhere tangent to the contact structure. The study of transverse and Legendrian knots is very important in both 3-dimensional contact geometry and low-dimensional topology. For a reference, see J. Etnyre's survey article [9].

In [4], Y. Eliashberg classified topologically trivial transverse knots in (R^3, ξ_0) up to transverse isotopy in terms of their self-linking number. In [6], Eliashberg–M. Fraser classified topologically trivial Legendrian knots in (R^3, ξ_0) up to Legendrian isotopy in terms of their Thurston–Bennequin invariant and rotation number. A topological knot type in (R^3, ξ_0) is called *transversely simple* (*Legendrian simple*, resp.) if transverse (Legendrian, resp.) knots in this type are determined by their self-linking number (Thurston–Bennequin invariant and rotation number, resp.). Therefore, in other words, Eliashberg and Eliashberg–Fraser showed that the trivial knot is transversely and Legendrian simple. Moreover recently, Etnyre and K. Honda proved that

torus knots and the figure-eight knot are both transversely and Legendrian simple ([8], [10]). For Legendrian non-simple knot types, Y. Chekanov and Eliashberg–H. Hofer independently discovered such examples ([3], [5], [7]). By computing a *contact homology invariant*, they succeeded in distinguishing a pair of Legendrian knots whose topological knot type, Thurston–Bennequin invariant and rotation number agree. Notice that their simplest topological knot type is a *twist knot* and a complete classification of a Legendrian twist knot except Legendrian trefoil and figure-eight knots is still not done. On the other hand, recently, J. Birman–W. Menasco and Etnyre–Honda, respectively, succeeded in finding examples of transversely non-simple knot type ([2], [11]). At writing this paper, however, it is not known whether a twist knot except the trefoil and figure-eight knots is transversely simple.

In this paper, we geometrically analyze transverse doubled and twist knots in (R^3, ξ_0) by using spanned clasp disks. As applications, we will estimate their self-linking number and furthermore we will show that in many cases, transverse twist knots with the maximal self-linking number are unique up to transverse isotopy.

2 Main Results

First we recall classical invariants of transverse and Legendrian knots in the standard contact 3-space (R^3, ξ_0) . Let v (h , resp.) be a non-singular vector field in ξ_0 (normal to ξ_0 , resp.) on R^3 .

Definition 1. The *self-linking number* $sl(T)$ of a transverse knot T in (R^3, ξ_0) is the linking number of T and a knot slightly pushed off T along v .

Definition 2. The *Thurston–Bennequin invariant* $tb(L)$ of a Legendrian knot L in (R^3, ξ_0) is the linking number of L and a knot slightly pushed off L along h . The *rotation number* $r(L)$ is the rotation number of L in the trivialization of ξ_0 given by v .

Notice that $sl(T)$ ($tb(L)$, resp.) is independent of choices of v (h , resp.) and an orientation of T (L , resp.) and that $r(L)$ is independent of a choice of v , but $r(L)$ changes its sign if the orientation of L is changed ([1]). In [1], D. Bennequin showed that $sl(T)$ and $tb(L) + |r(L)|$ are bounded by its negative Euler characteristic of Seifert surfaces for T and L , respectively.




Figure 1: $(J_{\pm}, n)$ -twist knots

A topological knot K in R^3 is a *doubled knot* of a companion knot K' if K is a satellite of K' obtained by an image of the knot J_+ or J_- in the solid torus $D^2 \times S^1$ featured in Figure 1, where the tubular neighbourhood of K' is replaced by the solid torus. A *twist knot* is a doubled knot whose companion knot is the trivial knot. A doubled knot is uniquely determined if the framing n of the image of $\{pt\} \times S^1, pt \in \partial D^2$ and the knot $J_{\pm}$ are chosen (n is an integer). In this paper, we call a doubled knot (twist knot, resp.) with the chosen framing n and J_+/J_- a $(J_+/J_-, n)$ -doubled knot ($(J_+/J_-, n)$ -twist knot, resp.). For example, let $+3_1, -3_1, 4_1$ denote the positive trefoil, negative trefoil, figure-eight knots, respectively. Then $+3_1$ is the $(J_+, -1)$ -twist knot, -3_1 is the $(J_-, 1)$ -twist knot, and 4_1 is both the $(J_+, 1)$ -twist knot and the $(J_-, -1)$ -twist knot.

For a topological knot type K , let $\bar{sl}(K)$ ($\bar{tb}(K)$, resp.) be the maximal value of the self-linking number (Thurston–Bennequin invariant, resp.) of all transverse (Legendrian, resp.) representative of K . For example, $\bar{sl}(O) = \bar{tb}(O) = -1$, $\bar{sl}(+3_1) = \bar{tb}(+3_1) = 1$, $\bar{sl}(-3_1) = -5$, $\bar{tb}(-3_1) = -6$, $\bar{sl}(4_1) = \bar{tb}(4_1) = -3$, where O denotes the trivial knot (for example, see [10]).

Theorem 1. Let K be a (J_+, n) -doubled knot of a companion knot K' . Suppose




Figure 2: A xz -projection of a transverse (J_+, n) -twist knot

$\bar{sl}(K) = 2\bar{tb}(K') - 2n + 1$. Further suppose Legendrian knots in the knot type K' with the maximal Thurston–Bennequin invariant are unique up to Legendrian isotopy. Then, transverse knots in the knot type K with the maximal self-linking number are unique up to transverse isotopy.

Corollary 1. Let K be a (J_+, n) -twist knot for $n \geq -1$. Then, transverse knots in the knot type K with the maximal self-linking number are unique up to transverse isotopy.

Proof of Corollary 1. By the Eliashberg’s theorem in [4], topologically trivial transverse knots in (R^3, ξ_0) with the maximal self-linking number -1 are unique. Therefore, by Theorem 1 it is sufficient to show that $\bar{sl}(K) = -2n - 1$. If $n \geq -1$, the lowest degree in the first variable of the HOMFLY polynomial of K is easily calculated to be $-2n - 1$. So we have $\bar{sl}(K) \leq -2n - 1$ by the HOMFLY bound of self-linking number ([12], [14]). Finally, let $\bar{K}$ be a transverse (J_+, n) -twist knot given by a generic projection on xz -plane featured in Figure 2. Note that the numbers of positive and negative crossings are 2 and $2n + 3$, respectively and the writhe of $\bar{K}$ is $-2n - 1$. Therefore, it follows that $sl(\bar{K}) = -2n - 1$. Hence the inequality is sharp.

Remark 1. The author does not know whether transverse knots in every twist knot type with the maximal self-linking number are unique up to transverse isotopy.

Theorem 2. Let K be a (J_+, n) -doubled knot of a companion knot K' . Then we




Figure 3: A clasp disk

have $\bar{sl}(K) \leq 2\bar{tb}(K') - 2n + 1$ and $\bar{tb}(K) \leq 2\bar{tb}(K') - 2n + 1$.

Corollary 2. Let K be a (J_+, n) -twist knot. Then we have $\bar{sl}(K) \leq -2n - 1$ and $\bar{tb}(K) \leq -2n - 1$.

Remark 2. As mentioned in the proof of Corollary 1, if $n \geq -1$, the inequality $\bar{sl}(K) \leq -2n - 1$ of Corollary 2 is sharp and also deduced from the HOMFLY bound. For a (J_+, n) -twist knot with $n \leq -2$, this is not sharp in general. In [15], L. Ng proved that the Kauffman bound of the maximal Thurston–Bennequin invariant for a two-bridge knot is sharp. Since a twist knot is a two-bridge knot, the inequality $\bar{tb}(K) \leq -2n - 1$ of Corollary 2 would be implied by that from the Kauffman bound.

3 Characteristic foliations on a clasp disk

Let K be an oriented doubled knot with companion K' in R^3 and let $\tilde{D}$ be a two-disk. Then there exists an immersion $f : \tilde{D} \rightarrow R^3$ such that $f|_{\partial\tilde{D}} : \partial\tilde{D} \rightarrow K$ is a diffeomorphism and $f^{-1}(\Omega)$ consists of two arcs $\tilde{\sigma}_1, \tilde{\sigma}_2$ joining a point in $\partial\tilde{D}$ and a point of $f^{-1}(K \cap f(Int(\tilde{D})))$ in $Int\tilde{D}$, where $\Omega = \{x \in f(\tilde{D}) \mid |f^{-1}(x)| \geq 2\}$. Put $D = f(\tilde{D})$. Throughout, we use the notation ∂D as $f(\partial\tilde{D})$ (ie., $\partial D = K$). Then D is an oriented immersed disk in R^3 such that the induced orientation of ∂D agrees with the orientation of K . We call D a *clasp disk* for K (Figure 3) and $\sigma = f(\tilde{\sigma}_1) = f(\tilde{\sigma}_2)$ the *clasp singularity*. Throughout this paper, for a $(J_+/J_-, n)$ -doubled knot, we only consider a naturally spanned clasp disk via J_+/J_- , respectively.

Further, suppose K is a transverse knot in (R^3, ξ_0) . Let D_{ξ_0} be the *characteristic foliation* of D , that is, the singular foliation defined by $TD \cap \xi_0$ on D . Remark that




Figure 4: A characteristic foliation

D_{ξ_0} is orientable. Throughout, we orient D_{ξ_0} in such a way that trajectories of D_{ξ_0} exit through ∂D .

Generically D_{ξ_0} satisfies the following:

- D is tangent to ξ_0 at a finite set Σ in $D - \partial D$ and all singular point in Σ is an *elliptic* point or a *hyperbolic* point according to the index (1 or -1 , respectively),
- there is no separatrix connection between hyperbolic points.

See Figure 4. In this paper, we call a clasp disk satisfying these conditions a *generic* clasp disk.

And we can distinguish between *positive* and *negative* points of Σ depending on whether the orientations of D and ξ_0 coincide at these points or not.

Let e_+ (e_- , resp.) and h_+ (h_- , resp.) be the number of positive (negative, resp.) elliptic and positive (negative, resp.) hyperbolic points in D_{ξ_0} , respectively. Then from D_{ξ_0} , we can compute the self-linking number of ∂D as follows.

Proposition 1 ([1, p. 133; Theorem 7]). $sl(\partial D)$ is equal to $\rho(D) - e_+ + e_- + h_+ - h_-$, where $\rho(D)$ is the linking number of ∂D and a knot obtained by shifting ∂D along the normal to D in R^3 .

Remark 3. $\rho(D)$ is ± 2 according to the sign of $J_{\pm}$.

In the next section, we will prove the following.

Theorem 3. For any transverse (J_+, n) -doubled knot K in (R^3, ξ_0) , there is a generic clasp disk D for K such that the clasp singularity σ of D is Legendrian and ξ_0 is transverse to D along σ . In particular, there is no singular point of the characteristic

foliation D_{ξ_0} along σ .

The goal of this section is to prove the following.

Theorem 4. Let K be a transverse doubled knot in (R^3, ξ_0) . Suppose that the self-linking number of K is maximal and that there is a generic clasp disk D' for K such that the clasp singularity σ' of D' is Legendrian and ξ_0 is transverse to D' along σ' . Then there exists a new generic clasp disk D for K satisfying the following:

- the clasp singularity σ of D is Legendrian and ξ_0 is transverse to D along σ ,
- $e_- = h_+ = 0$ in D_{ξ_0} ,
- there is a Legendrian embedded circle C in D which contains σ and all positive elliptic and negative hyperbolic point in D_{ξ_0} alternatively along $C - \sigma$.

Let K be a transverse doubled knot in (R^3, ξ_0) and D be a generic clasp disk for K . Suppose that the clasp singularity σ of D is Legendrian and ξ_0 is transverse to D along σ .

First, as fully discussed in [4], we will modify D relative to both ∂D and σ so that $e_- = h_+ = 0$ in D_{ξ_0} .

The next lemma due to E. Giroux is a basic tool of a modification of D_{ξ_0} which allows us to cancel or create the pairs of an elliptic point and a hyperbolic point of the same sign in D_{ξ_0} .

Lemma 1 (Elimination lemma [4], [13]). Let $p, q \in D - \sigma$ be elliptic and hyperbolic points of D_{ξ_0} of the same sign. Let γ be a trajectory of D_{ξ_0} which joins p and q and is disjoint from σ . Then there exists a C^0 -small isotopy $f_t : \tilde{D} \rightarrow R^3, t \in [0, 1]$, which starts with $f_0 = f$, is fixed on γ and outside of a neighbourhood $U \supset \gamma$ and such that the characteristic foliation of $f_1(\tilde{D})$ has no singularity in $U \cap f_1(\tilde{D})$.

We first eliminate all negative elliptic point in D_{ξ_0} . Let p be a negative elliptic point in D_{ξ_0} . If there is a negative hyperbolic point connected with p by a trajectory in D_{ξ_0} , then we cancel the pair by Lemma 1. If otherwise, as in the proof of Theorem 4.1.1 in [4], after creating the pairs of positive elliptic and positive hyperbolic points, if necessary, we have a limit cycle in D_{ξ_0} which is disjoint from both ∂D and σ . This contradicts the Bennequin inequality.

Next, we eliminate all positive hyperbolic point in D_{ξ_0} . Let q be a positive hy-




Figure 5:

perbolic point in D_{ξ_0} . Then by our assumptions, a stable separatrix of q necessarily connects q with a positive elliptic point. Again we cancel the pair by Lemma 1.

It follows that after a small isotopy of D relative to both ∂D and σ , we can assume that there is a map α from an abstract finite tree G to R^3 satisfying:

- $e_- = h_+ = 0$ in D_{ξ_0} ,
- $\alpha(G)$ contains all singular point in D_{ξ_0} ,
- α is injective,
- the image of α of all vertex of G equals the set of all positive elliptic point in D_{ξ_0} ,
- the restriction of α to each edge of G is Legendrian and the each image contains exactly one interior singular point which is negative hyperbolic.

See Figure 5.

Next we recall *stabilizations* and *bypasses* ([8], [10]).

Definition 3. A transverse knot T is a *single stabilization* of a transverse knot T' if $T = \beta \cup A, T' = \beta \cup A'$ and $A \cup A'$ cobounds a disk with only one positive elliptic and one negative hyperbolic points and $sl(T) = sl(T') - 2$ (Figure 6). The disk bounded




Figure 6: A transverse bypass

by $A \cup A'$ is called a *transverse bypass* and T' is called a *single destabilization* of T .

Let D_{ξ_0} be as above.

Now after doing single destabilizations on transverse bypasses on D_{ξ_0} disjoint from the clasp singularity σ , if necessary, we have a generic clasp disk D' such that after a small modification of D'_{ξ_0} , there is a Legendrian embedded arc δ in D' which satisfies the following:

- δ contains all positive elliptic and negative hyperbolic point alternatively along it in D'_{ξ_0} ,
- δ is disjoint from the clasp singularity σ' of D' ,
- $\partial\delta$ is two positive elliptic points in D'_{ξ_0} .

Further, by the genericity of D'_{ξ_0} , we can assume that σ' is not included in an unstable trajectory from a negative hyperbolic point. See Figure 7.

Then finally, suppose that K has the maximal self-linking number. Since a single destabilization increases the self-linking number by two, it follows that in the above D'_{ξ_0} , δ can be extended by unstable trajectories from $\partial\delta$ to σ' so that the union of the extended δ and σ' is a simple closed curve in D' . Because if otherwise, we can do more single destabilization.

After smoothing the corners of the simple closed curve, we have a desired Legendrian embedded circle C in D' as in the statement of Theorem 4. This completes the proof of Theorem 4.




Figure 7:

Remark 4. In Theorem 4, if K is a $(J_{\pm}, n)$ -doubled knot, $tb(C) = n - e_+ + 1 = n - h_-$.

4 Proof of Theorem 3

For the proof of Theorem 3, we need a little knowledge of Legendrian knot theory for the standard contact 3-space (R^3, ξ_0) and convex surface theory as in [10], [13] and [16].

Let K be a transverse (J_+, n) -doubled knot.

First, take a *Legendrian push off* L of K such that L is C^0 -close to K and L is a closed leaf on the boundary of a standard neighbourhood of K ([4], [9]). We remark that a Legendrian isotopy of L induces a transverse isotopy of K ([4], [9]). Let E be a clasp disk naturally spanned by the Legendrian (J_+, n) -doubled knot L and let c be a simple closed curve on E such that c contains the clasp singularity of E and that the topological knot type of c is the companion of K .

Then it is not hard to see that there are a topologically standard solid torus V embedded in (R^3, ξ_0) and meridian disks d_1, d_2 of V which satisfy the following (Figure 8):

- $E \subset \text{Int}V$,
- E intersects d_i as an arc ($i = 1, 2$),
- L intersects d_i geometrically twice and algebraically zero ($i = 1, 2$),
- a 3-ball B cut by d_1 and d_2 in V contains a neighbourhood of the clasp singularity




Figure 8:




Figure 9: A front

of E ,

- the intersection of c and B is an unknotted arc properly embedded in B ,
- ∂V is a convex surface whose dividing set is two parallel simple closed curves and their slopes are a negative integer,
- ∂d_i is Legendrian and d_i is a convex disk whose dividing set is an arc ($i = 1, 2$).

Then as discussed in [16], by a contact isotopy of (R^3, ξ_0) we can regard V as a “thin” tubular neighbourhood of a Legendrian unknot in (R^3, ξ_0) . Without loss of generality, we consider L is given by a *front* in xz -plane in (R^3, ξ_0) . So we can assume that B, d_i and $L|_B$ are corresponding to a square, its vertical edge and a front as in Figure 9, respectively ($i = 1, 2$).

By using Legendrian Reidemeister moves in the square, if necessary, we can find a Legendrian clasp part for L featured in Figure 10. Then it is easy to see that there is a clasp disk D_L for L such that along the interior of its clasp singularity σ_L , there is no tangency of D_L to ξ_0 , where σ_L is indicated by the dotted segment in Figure 10.

From D_L , we can easily make the desired clasp disk for K . This completes the




Figure 10: Legendrian clasps

proof.

Remark 5. For a transverse (J_-, n) -doubled knot, Theorem 3 does not hold in general. For instance, suppose a transverse $(J_-, 0)$ -twist knot K with the maximal self-linking number spans such a clasp disk D . Then by Theorem 4, we can assume that $e_- = h_+ = 0$ and $e_+ = h_- + 1$ in D_{ξ_0} . Therefore by Proposition 1, we have $sl(\partial D) = -1 - 2e_+$. But since $K = \partial D$ is topologically trivial and $sl(K) = -1$, it follows that $e_+ = 0$, a contradiction.

5 Proof of Theorem 1 and Theorem 2

We are ready to prove Theorem 1. Let K_1 and K_2 be transverse (J_+, n) -doubled knots. Suppose that K_1 is equivalent to K_2 as a topological knot K and that both K_1 and K_2 have the maximal self-linking number. Without loss of generality, we assume both K_1 and K_2 have the same companion knot K' . By Theorem 3 and Theorem 4, there exist a generic clasp disk D_i for K_i and a Legendrian circle C_i in D_i as in the statement of Theorem 4 ($i = 1, 2$). Remark that $D_{1\xi_0}$ is diffeomorphic to $D_{2\xi_0}$. We orient C_1 and C_2 in the same way along D_1 and D_2 . C_1 is equivalent to C_2 as the companion knot of K_i . Since $sl(K_1) = sl(K_2)$, we have $tb(C_1) = tb(C_2)$ by Proposition 1 and Remark 4.

Now suppose $\bar{sl}(K) = 1 - 2n + 2\bar{tb}(K')$. Then by Proposition 1, Remark 3 and Remark 4, we have $tb(C_i) = \bar{tb}(C_i)$ ($i = 1, 2$).

Further suppose Legendrian knots in the knot type K' with the maximal Thurston–Bennequin number are unique. Then, it follows that C_1 and C_2 are Legendrian isotopic. So there exists an orientation preserving diffeotopy $g_t : R^3 \rightarrow R^3$ such that $g_0 = \text{id}$, g_t is a contactomorphism of ξ_0 for each t , and $g_{1*}(C_1) = C_2$ with its orientations. Without loss of generality, we can further assume $g_1(\sigma_1) = \sigma_2$, where σ_1, σ_2 are the

clasp singularities of D_1, D_2 . Then, using g_t and D_1, D_2 , it is not hard to see K_1 and K_2 are transversely isotopic. This completes the proof of Theorem 1.

Proof of Theorem 2. Let K be a transverse (J_+, n) -doubled knot with companion K' . Suppose $sl(K) = \bar{sl}(K)$. Then by Theorem 3 and Theorem 4, there exist a clasp disk D for K and a Legendrian circle C in D as in the statement of Theorem 4. By Proposition 1 and Remark 4, we have $2tb(C) - 2n = \bar{sl}(K) - 1$. Since C is topologically equivalent to K' , it follows that $\bar{sl}(K) \leq 2\bar{tb}(K') - 2n + 1$.

And let L be a Legendrian knot which is topologically equivalent to K . Suppose $tb(L) = \bar{tb}(K)$. Since $sl(T_{\pm}(L)) = tb(L) \mp r(L)$, where $T_+(L), T_-(L)$ are positive and negative transverse push offs of L , we also have $\bar{tb}(K) \leq 2\bar{tb}(K') - 2n + 1$.

References

- [1] D. Bennequin, *Entrelacements et équations de Pfaff*, Astérisque, **107-108** (1983) 87–161.
- [2] J. Birman–W. Menasco, *Stabilization in the braid groups II: On transversal simplicity of transversal knots*, preprint.
- [3] Y. Chekanov, *Differential algebra of Legendrian links*, Invent. Math., **150** (2002) 441–483.
- [4] Y. Eliashberg, *Legendrian and transversal knots in tight contact 3-manifolds*, Topological Methods in Modern Mathematics, (1993) 171–193.
- [5] Y. Eliashberg, *Invariants in contact topology*, Proceedings of the ICM (Berlin, 1998), Doc. Math. **2** (1998) 327–338.
- [6] Y. Eliashberg–M. Fraser, *Classification of topologically trivial Legendrian knots*, Geometry, topology, and dynamics (Montreal, PQ, 1995), CRM Proc. Lecture Notes, AMS, **15** (1998) 17–51.
- [7] J. Epstein–D. Fuchs–M. Meyer, *Chekanov-Eliashberg invariants and transverse approximations of Legendrian knots*, Pacific J. Math., **201** (2001) 89–106.
- [8] J. Etnyre, *Transversal torus knots*, Geometry and Topology, **3** (1999) 253–268.
- [9] J. Etnyre, *Legendrian and transversal knots*, to appear in the Handbook of Knot Theory.

- [10] J. Etnyre–K. Honda, *Knots and Contact geometry I: torus knots and the figure eight*, J. Symplectic Geometry, **1** (2001) 63–120.
- [11] J. Etnyre–K. Honda, *Cabling and transverse simplicity*, preprint.
- [12] D. Fuchs–S. Tabachnikov, *Invariants of Legendrian and transverse knots in the standard contact space*, Topology, **66** (1997) 1025–1053.
- [13] E. Giroux, *Convexité en topologie de contact*, Comment. Math. Helvetici, **66** (1991) 637–677.
- [14] V. Goryunov–J. Hill, *A Bennequin number estimate for transverse knots*, ‘Singularity Theory. Proceedings of the Conference dedicated to C.T.C.Wall’s 60th birthday’, London Mathematical Society Lecture Notes Series, **263** (1999) 265–280.
- [15] L. Ng, *Maximal Thurston-Bennequin Number of Two-Bridge Links*, Algebraic and Geometric Topology, **1** (2001) 427–434.
- [16] I. Torisu, *On the additivity of the Thurston–Bennequin invariant of Legendrian knots*, Pacific J. Math., **210** (2003) 359–365.